

Properties of 2×2 h -deformed quantum (super)matrices*

Yun Li[†] and Sicong Jing

*Department of Modern Physics,
University of Science and Technology of China,
Hefei, Anhui 230026, P.R.China*

February 1, 2008

Abstract

We investigate the h -deformed quantum (super)group of 2×2 matrices and use a kind of contraction procedure to prove that the n -th power of this deformed quantum (super)matrix is quantum (super)matrix with the deformation parameter nh .

keywords: h -deformed, quantum group, quantum matrix

There are two distinct quantizations for general Lie group. One is the well-known Drinfeld-Jimbo q -deformed quantum group and the other is the so-called Jordanian h -deformed one. For group $GL(2)$ there exist only such two deformation (up to isomorphism) with a central quantum determinant: $GL_q(2)$ and $GL_h(2)$ [1]. One can obtain these quantum groups by deforming the coordinates of a linear plane to be noncommutative objects. In detail, the deformed quantum groups act on the q -plane, with the relation $xy = qyx$ and the h -plane, with the relation $xy - yx = hy^2$, respectively. The q -deformed quantum group has been intensively studied[2] because it is closely related with the solution of Yang-Baxter equation and the theory of braids. In [3], it is found that the deformation parameter of h -deformed quantum group is naturally dimensional (like the known k -deformation). This is quite intriguing for possible physical applications. Recently geometrical structure of the h -plane has been thoroughly discussed[4] and quantum mechanics based on the plane was constructed[5]. These work are helpful to understand h -deformed quantum group.

In [6], q -deformed quantum group $GL_q(2)$ was investigated and it was shown that the n -th power of a q -deformed quantum matrix corresponds to the n -th power of the deformation parameter q . Also the same property of q -deformed quantum supergroup $GL_q(1|1)$ was discussed in [7]. And in [1], it was pointed

*The project supported by National Natural Science Foundation of China under Grant No. 10075042

[†]Email: yunlee@mail.ustc.edu.cn

out that the n -th power of a h -deformed quantum matrix is also quantum matrix with the deformation parameter nh . In [8], it was found that the h -deformed quantum group can be obtained from the q -deformed quantum Lie group through a singular limit $q \rightarrow 1$ of a linear transformation. This method is called as the contraction procedure.

In this paper we use the contraction method to obtain the same result pointed out in [1]. We also extend the discussion to deformed quantum supergroup and show that the n -th power of a h -deformed quantum supermatrix is also quantum supermatrix with the deformation parameter nh . Not only do we use the singular linear transformation between the two kinds of quantum (super)groups to obtain the results, but also we use the similarity transformation between the corresponding R -matrices for different quantum groups to obtain them.

To begin with we define the q -deformed quantum matrix as

$$M' = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}. \quad (1)$$

Throughout this paper we denote q -deformed objects by primed quantities. The unprimed ones represent transformed deformed objects. The Manin quantum plane satisfying the commutation relation $xy = qyx$ can be considered covariant with respect to the action of the quantum group $GL_q(2)$. $M' \in GL_q(2)$ means the following commutation relations are fulfilled

$$\begin{aligned} a'c' &= qc'a', & b'd' &= qd'b', & [a', d'] &= (q - q^{-1})b'c', \\ a'b' &= qb'a', & c'd' &= qd'c', & c'b' &= b'c'. \end{aligned} \quad (2)$$

Here $[,]$ stands for the commutator. The quantum determinant D' which is defined as

$$D' = \det_q M' \equiv a'd' - qc'b' \quad (3)$$

is central. One can obtain h -deformed quantum group $GL_h(2)$ with the following transformation

$$M' = gMg^{-1} \quad (4)$$

where

$$g = \begin{pmatrix} 1 & \frac{h}{q-1} \\ 0 & 1 \end{pmatrix}. \quad (5)$$

A simple calculation shows that the q -deformed matrix elements and the corresponding transformed ones fulfil the following relations:

$$\begin{aligned}
a' &= a + \frac{h}{q-1}c, & b' &= b + \frac{h}{q-1}(d-a) - \frac{h^2}{(q-1)^2}c, \\
c' &= c, & d' &= d - \frac{h}{q-1}c.
\end{aligned} \tag{6}$$

Substituting the above relations into the q -commutation relations(2) and taking the limit of $q \rightarrow 1$, one can find that the entries of the transformed quantum matrix M satisfy the following commutation relations of the $GL_h(2)$

$$\begin{aligned}
[a, c] &= hc^2, & [d, b] &= h(D_h - d^2), & [a, d] &= hdc - hac, \\
[d, c] &= hc^2, & [b, c] &= hac + hcd, & [b, a] &= h(a^2 - D_h).
\end{aligned} \tag{7}$$

Here the corresponding h -deformed determinant is

$$D_h = \det_h M \equiv ad - cb - hcd \tag{8}$$

which is also central. The h -deformed quantum plane satisfying the relation $xy = yx + hy^2$ can also be considered covariant with respect to the action of the above quantum group $GL_h(2)$.

For n -th power of M' , from (4), we obtain

$$\begin{aligned}
M'^n &= (gMg^{-1})^n \\
&= gM^n g^{-1}.
\end{aligned} \tag{9}$$

If we denote M'^n, M^n as $\begin{pmatrix} a'_n & b'_n \\ c'_n & d'_n \end{pmatrix}$ and $\begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}$, respectively, the relations between them can be obtained in a similar way to get Eq.(6). They are given by

$$\begin{aligned}
a'_n &= a_n + \frac{h}{q-1}c_n, & b'_n &= b_n + \frac{h}{q-1}(d_n - a_n) - \frac{h^2}{(q-1)^2}c_n, \\
c'_n &= c_n, & d'_n &= d_n - \frac{h}{q-1}c_n.
\end{aligned} \tag{10}$$

But because M'^n belongs to $GL_{q^n}(2)$ [6], the matrix elements of M'^n will satisfy the commutation relations(2) only with the deformed parameter q replaced by q^n . So we can have the following relations

$$\begin{aligned}
a'_n c'_n &= q^n c'_n a'_n, & b'_n d'_n &= q^n d'_n b'_n, & [a'_n, d'_n] &= (q^n - q^{-n}) b'_n c'_n, \\
a'_n b'_n &= q^n b'_n a'_n, & c'_n d'_n &= q^n d'_n c'_n, & c'_n b'_n &= b'_n c'_n.
\end{aligned} \tag{11}$$

Following the same procedure as we obtain relations(7) and with the limit identity

$$\lim_{q \rightarrow 1} \frac{q^n - 1}{q - 1} = n , \quad (12)$$

we can have the following commutation relations of the $GL_{nh}(2)$ given by

$$\begin{aligned} [a_n, c_n] &= nhc_n^2, & [d_n, b_n] &= nh(D_{nh} - d_n^2), \\ [a_n, d_n] &= nhd_nc_n - nha_nc_n, & [d_n, c_n] &= nhc_n^2, \\ [b_n, c_n] &= nha_nc_n + nhc_nd_n, & [b_n, a_n] &= nh(a_n^2 - D_{nh}) \end{aligned} \quad (13)$$

where

$$D_{nh} = \det_h M^n \equiv a_nd_n - c_nb_n - nhc_nd_n, \quad (14)$$

which mean that M^n does belong to the h -deformed quantum group $GL_{nh}(2)$.

In fact, by using the R -matrix for q -deformation[6]:

$$R' = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & q & 1 - q^2 & 0 \\ 0 & 0 & q & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (15)$$

one can obtain the commutation relations(2) between the elements of q -deformed matrix M' through the following equation

$$R' M'_1 M'_2 = M'_2 M'_1 R', \quad (16)$$

where M'_1, M'_2 are the graded matrices with the following matrix elements respectively

$$\begin{aligned} (M'_1)_{cd}^{ab} &= (M' \otimes I)_{cd}^{ab} = M'_c{}^a \delta_d^b, \\ (M'_2)_{cd}^{ab} &= (I \otimes M')_{cd}^{ab} = M_d{}^b \delta_c^a. \end{aligned} \quad (17)$$

And it is clear that the transformation between the deformed quantum groups $M' = gMg^{-1}$ leads to the similarity transformation between the corresponding R -matrices for different quantum groups:

$$R = (g \otimes g)^{-1} R' (g \otimes g). \quad (18)$$

Here product $\otimes$ acts in the same manner as M'_1 and M'_2 expressions(17). Through such transformation the R -matrix for the h -deformation can be obtained as the

following

$$R_h \equiv \lim_{q \rightarrow 1} R = \begin{pmatrix} 1 & -h & h & h^2 \\ 0 & 1 & 0 & -h \\ 0 & 0 & 1 & h \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (19)$$

Applying Eq.(4) and Eq.(18), one can obtain the following equation in the limit of $q \rightarrow 1$

$$R_h M_1 M_2 = M_2 M_1 R_h. \quad (20)$$

So the h -commutation relations(7) can be derived from the above equation again. As a common sense, the R -matrix for q -deformation in the quantum group $GL_{q^n}(2)$ is given by

$$R'_n = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & q^n & 1 - q^{2n} & 0 \\ 0 & 0 & q^n & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (21)$$

Because of the transformation(9), we also have

$$R_n = (g \otimes g)^{-1} R'_n (g \otimes g). \quad (22)$$

So with the limit identity(12), we can obtain the R -matrix for h -deformation in the quantum group $GL_{nh}(2)$ following by

$$R_{nh} \equiv \lim_{q \rightarrow 1} R_n = \begin{pmatrix} 1 & -nh & nh & (nh)^2 \\ 0 & 1 & 0 & -nh \\ 0 & 0 & 1 & nh \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (23)$$

Also from Eq.(9) and Eq.(22), we derive the following equation in the limit of $q \rightarrow 1$

$$R_{nh} M_1^n M_2^n = M_2^n M_1^n R_{nh} \quad (24)$$

and the commutation relations Eq.(13) of the quantum group $GL_{nh}(2)$ are obtained through this equation and it shows that M^n definitely belongs to $GL_{nh}(2)$.

Next step, we will extend our discussion to quantum supergroup. We denote supermatrix quantity as $\tilde{Q}$ with Q representing the quantity for the common deformed group. Returning to the Manin quantum superplane satisfying the relation $x\theta = q\theta x$, $\theta^2 = 0$, there exists the R -matrix for the quantum supergroup $GL_q(1 | 1)$ acting on the superplane[9], where

$$\tilde{R}' = \begin{pmatrix} q & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & q - q^{-1} & 1 & 0 \\ 0 & 0 & 0 & q^{-1} \end{pmatrix}. \quad (25)$$

The h -deformed quantum superplane satisfying the relation $x\theta = \theta x + hx^2$, $\theta^2 = -h\theta x$ can also be obtained through the singular transformation in the limit $q \rightarrow 1$. Conventionally, here we take the transfer matrix g as $\begin{pmatrix} 1 & 0 \\ \frac{h}{q-1} & 1 \end{pmatrix}$.

Observing that for h -deformed quantum supergroup the deformation parameter must be a dual odd (Grassmann) number satisfying $h^2 = 0$ [10], one can obtain the corresponding R -matrix for the quantum supergroup $GL_h(1 | 1)$ acting on the h -deformed quantum superplane through the similarity transformation in Eq.(18):

$$\tilde{R}_h \equiv \lim_{q \rightarrow 1} \tilde{R} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -h & 1 & 0 & 0 \\ h & 0 & 1 & 0 \\ 0 & h & h & 1 \end{pmatrix}. \quad (26)$$

Here, the only difference is the grading product $\otimes$. For quantum supergroup, the product acts in the following manner

$$\begin{aligned} (\tilde{M}'_1)_{cd}^{ab} &= (\tilde{M}' \otimes I)_{cd}^{ab} = (-1)^{c(b+d)} \tilde{M}'_c{}^a \delta_d^b, \\ (\tilde{M}'_2)_{cd}^{ab} &= (I \otimes \tilde{M}')_{cd}^{ab} = (-1)^{a(b+d)} \tilde{M}'_d{}^b \delta_c^a. \end{aligned} \quad (27)$$

From the same relation as Eq.(20), one can obtain the commutation relations of quantum supergroup $GL_h(1 | 1)$, which is given by

$$\begin{aligned} ab &= ba, & ac &= ca + h(a^2 + cb - da) \\ bd &= db, & cd &= dc + h(d^2 - cb - da) \\ b^2 &= 0, & c^2 &= h(dc - ca) \\ bc &= -cb - h(ba - db), & ad &= da + h(ba - db). \end{aligned} \quad (28)$$

The n -th power of supermatrix $\tilde{M}'$ belongs to quantum supergroup $GL_{q^n}(1 | 1)$ and R -matrix for $GL_{q^n}(1 | 1)$ is given by

$$\tilde{R}'_n = \begin{pmatrix} q^n & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & q^n - q^{-n} & 1 & 0 \\ 0 & 0 & 0 & q^{-n} \end{pmatrix}. \quad (29)$$

So through the transformation like Eq.(22) and with the limit of $q \rightarrow 1$, we can have the R -matrix for quantum supergroup $GL_{nh}(1 | 1)$ as following

$$\tilde{R}_{nh} \equiv \lim_{q \rightarrow 1} \tilde{R} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -nh & 1 & 0 & 0 \\ nh & 0 & 1 & 0 \\ 0 & nh & nh & 1 \end{pmatrix} \quad (30)$$

Also following the same procedure as we obtain Eq.(24) we have the relation for the n -th power of quantum supermatrix $\widetilde{M}^n$:

$$\widetilde{R}_{nh} \widetilde{M}_1^n \widetilde{M}_2^n = \widetilde{M}_2^n \widetilde{M}_1^n \widetilde{R}_{nh} , \quad (31)$$

which shows that the entries of the transformed quantum matrix $\widetilde{M}^n$ fulfil the commutation relations of the quantum supergroup $GL_{nh}(1 | 1)$ as following

$$\begin{aligned} ab &= ba, & ac &= ca + nh(a^2 + cb - da) \\ bd &= db, & cd &= dc + nh(d^2 - cb - da) \\ b^2 &= 0, & c^2 &= nh(dc - ca) \\ bc &= -cb - nh(ba - db), & ad &= da + nh(ba - db) . \end{aligned} \quad (32)$$

Hence, we conclude that $\widetilde{M}^n$ belongs to $GL_{nh}(1 | 1)$.

In summary, we show that the n -th power of h -deformed quantum matrix is quantum matrix with the deformation parameter nh and the same is also true for the n -th power of h -deformed quantum supermatrix.

References

- [1] Kupershmidt, B A. (1992). *J. Phys. A: Math. Gen.* **25** L1239
- [2] Drinfeld, V G. (1986). *Quantum groups*, Proceeding of the International Congress of Mathematicians, Berkley, Vol **1**, 798
Faddeev, L D., Reshetikhin, N Y., and Takhtajan, L A. (1989) *Algebra Anal* **1**, 178
- [3] Lukierski, J., Ruegg, H., and Tolostoy, V N., (1995). ICTP preprint
- [4] Cho, S., Madore, J., and Park, K S., (1998). *J. Phys. A: Math. Gen.* **31** 2639
- [5] Cho, S. (1999). *J. Phys. A: Math. Gen.* **32** 2091
- [6] Vokos, S P., Zumino, B., and Wess, J., (1990). *Z. Phys. C: Particles and Fields* **48** 65
Corrigan, E., Fairlie, D B., and Fletcher, P. (1990). *J. Math. Phys.* **31** 776
- [7] Schwenk, J., Schmidke, W B., and Vokos, S P. (1990). *Z. Phys. C: Particles and Fields* **46** 643
- [8] Aghamohammadi, A., Khorrami, M., and Shariati, A. (1995). *J. Phys. A: Math. Gen.* **28** L225
- [9] Schmidke, W B., Vokos, S P., and Zumino, B. (1990) *Z. Phys. C: Particles and Fields* **48**, 249
- [10] Dabrowski, L., and Parashar, P. (1996). *Lett. Math. Phys.* **38** 331